



Assessing Benefits and Harms of Opioid Therapy

The National Epidemic

- The United States is in the midst of an epidemic of prescription opioid overdose deaths, which killed more than 14,000 people in 2014 alone.
- Since 1999, sales of prescription opioids—and related overdose deaths—have quadrupled.
- Since 1999, there have been more than 165,000 deaths from overdose related to prescription opioids.

The Utah Epidemic

- Utah has the seventh highest drug overdose rate in the U.S.
- From 2013 to 2015, Utah ranked 7th highest in the nation for drug overdose deaths.¹
- In 2015, 24 individuals (residents and non-residents) died every month from a prescription opioid overdose in Utah.²

Guidance for Prescribing Opioids

The Utah Clinical Guidelines on Prescribing Opioids for Treatment of Pain³ provides up-to-date guidance on prescribing and weighing the risks and benefits of opioids.

- Before starting and periodically during opioid therapy, discuss the known risks and realistic benefits of opioids.
- Also discuss provider and patient responsibilities for managing therapy.
- Within 1-4 weeks of starting opioid therapy, and at least every three months, evaluate benefits and harms with the patient.

Assess benefits of opioid therapy

Assess your patient's pain and function regularly. A 30 percent improvement in pain and function is considered clinically meaningful. Discuss patient-centered goals and improvements in function (such as returning to work and recreational activities) and assess pain using validated instruments such as the 3-item (PEG) Assessment Scale:

1. What number best describes your pain on average in the past week? (from 0=no pain to 10=pain as bad as you can imagine)
2. What number best describes how, during the past week, pain has interfered with your enjoyment of life? (from 0=does not interfere to 10=completely interferes)
3. What number best describes how, during the past week, pain has interfered with your general activity? (from 0=does not interfere to 10=completely interferes)

If your patient does not have a 30 percent improvement in pain and function, consider reducing dose or tapering and discontinuing opioids. Continue opioids only as a careful decision by you and your patient when improvements in both pain and function outweigh the harms.

STOP THE OPIIDEMIC

Assess Harms of Opioid Therapy

Long-term opioid therapy can cause harms ranging in severity from constipation and nausea to opioid use disorder and overdose death. Certain factors can increase these risks, and it is important to assess and follow up regularly to reduce potential harms.

- 1. ASSESS.** Evaluate for factors that could increase your patient's risk for harm from opioid therapy such as:
 - Personal or family history of substance use disorder
 - Anxiety or depression
 - Pregnancy
 - Age 65 or older
 - COPD or other underlying respiratory conditions
 - Renal or hepatic insufficiency
- 2. CHECK.** Consider urine drug testing for other prescription or illicit drugs and check your state's prescription drug monitoring program (PDMP) for:
 - Possible drug interactions (such as benzodiazepines)
 - High opioid dosage (≥ 50 MME/day)
 - Obtaining opioids from multiple providers
- 3. DISCUSS.** Ask your patient about concerns and determine any harms they may be experiencing such as:
 - Nausea or constipation
 - Feeling sedated or confused
 - Breathing interruptions during sleep
 - Taking or craving more opioids than prescribed or difficulty controlling use
- 4. OBSERVE.** Look for early warning signs for overdose risk such as:
 - Confusion
 - Sedation
 - Slurred speech
 - Abnormal gait

If harms outweigh any experienced benefits, work with your patient to reduce dose, or taper and discontinue opioids and optimize nonopioid approaches to pain management.

Tapering and Discontinuing Opioid Therapy

Symptoms of opioid withdrawal may include drug craving, anxiety, insomnia, abdominal pain, vomiting, diarrhea and tremors. Tapering plans should be individualized. However, in general:

- 1. Go Slow:** To minimize symptoms of opioid withdrawal, decrease 10 percent of the original dose per week. Some patients who have taken opioids for a long time might find slower tapers easier (e.g., 10 percent of the original dosage per month).
- 2. Consult:** Work with appropriate specialists as needed—especially for those at risk of harm from withdrawal such as pregnant patients and those with opioid use disorder.
- 3. Support:** During the taper, ensure patients receive psychosocial support for anxiety. If needed, work with mental health providers and offer or arrange for treatment of opioid use disorder.

Improving the way opioids are prescribed can ensure patients have access to safer, more effective chronic pain treatment while reducing the number of people who misuse, abuse or overdose from these drugs.

References

1. U.S. Centers for Disease Control and Prevention, Web-based Injury and Statistics.
2. Utah Department of Health Violence and Injury Prevention Program, Prescription Pain Medication Program Database.
3. Recommendations do not apply to pain management in the context of active cancer treatment, palliative care, and end-of-life care.